

Scientific Reports in Medicine

The relationship the mobbing situation of healthcare workers and their conflict action styles

Mobbing in Healthcare Workers

Merve Eserler¹, Nergiz Sevinç², ErKay Nacar³, Ali Ayberk Arıcan⁴

DOI 10.37609/srinmed.43

Abstract: **Objective:** This study aims to examine whether there is a relationship between exposure to mobbing behaviors and conflict action styles among healthcare workers in Karabük Province.

Material and Method: This is a descriptive and cross-sectional study conducted with 225 healthcare workers in Karabük between May 15, 2023, and August 15, 2023. The data were collected using the Sociodemographic Information Questionnaire, the Mobbing Scale, and the Conflict Action Styles Scale.

Results: The participants' average score on the mobbing scale was found to be 82.15 ± 44.10 . Among the sub-dimensions of the mobbing scale, the highest score was obtained in the "relationships with colleagues" sub-dimension. Participants scored the highest in the "facilitating approach" sub-dimension of the Conflict Action Styles Scale. A significant difference was found between healthcare workers' exposure to mobbing and their marital status, profession, workplace, weekly working hours, job satisfaction, and smoking status. Additionally, significant differences were found between conflict action styles and gender, education, profession, weekly working hours, and job satisfaction.

Conclusion: Healthcare professionals are at risk of experiencing mobbing. This is an important issue that needs attention in terms of employee satisfaction and quality of life. Emphasizing mobbing and conflict action styles in training, implementing more comprehensive legal regulations, and improving working conditions are considered to be beneficial measures.

Keywords: Healthcare workers, Mobbing, Conflict action styles.

¹Karabük Education and Research Hospital, Karabük, Türkiye
Email: merveeserler@gmail.com
ORCID iD: 0009-0009-6205-581X

²Karabük University Faculty of Medicine, Department of Public Health, Karabük, Türkiye
Email: mdesvnc@gmail.com
ORCID iD: 0000-0003-4763-1902

³Karabük University Faculty of Medicine, Department of Public Health, Karabük, Türkiye
Email: erkaynacar@karabuk.edu.tr
ORCID iD: 0000-0002-7046-4551

⁴Karabük University Faculty of Medicine, Department of Public Health, Karabük, Türkiye
Email: aliayerka@gmail.com
ORCID iD: 0009-0002-3281-1783

Received: 2024-12-25
Accepted: 2025-03-02

3023-8226 / Copyright © 2024 by Akademisyen Publishing. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

INTRODUCTION

Mobbing is defined as psychological harassment, violence, intimidation, pressure, and similar behaviors carried out by one or more individuals in the same work environment towards an employee (1). It is described as any form of abusive, repetitive, and systematic behavior that jeopardizes an individual's dignity, physical, or mental integrity (2). The word "mobbing" is derived from the Latin term "mobile vulgus," which means an unstable crowd, originating from the root "Mob." In the dictionary, "Mob" is defined as "a disorderly crowd that engages in illegal violence" or "bullying" (3, 4).

The concept of mobbing was first scientifically introduced into the workplace context by Swedish psychiatrist Heinz Leymann in the 1980s. Leymann coined the term to describe a specific form of bullying towards employees in the workplace (5). He also used the term "psychological terrorism" in relation to mobbing in workplaces (6). Mobbing has become a recognized international workplace issue in the literature. This issue is evident in any country and culture (7). Mobbing can occur in both the private and public sectors, and every individual in the workforce is a potential victim of mobbing (8).

All employees in the healthcare sector can be subjected to mobbing behaviors. Factors contributing to mobbing in the healthcare sector include the need for various professional groups to work together, continuous work demands, stressful work environments, insufficient wages, bureaucratic barriers, lack of medical facilities, and unclear job descriptions (9). Due to the matrix organizational structure in hospitals, communication problems between employees are frequently encountered. Insufficient communication leads to misunderstandings and problems among employees (10). A study conducted internationally concluded that healthcare workers in hospitals, due to their unique structures, are exposed to mobbing behaviors 16 times more than those working in other service sectors (11). A retrospective study on 60 physician suicides in Italy over the past decade

found that 20% of them were related to work problems involving mobbing (12).

Conflict is described as a situation where one person's demands, expectations, and interests are opposed, resisted, or differentiated by another group or individuals, creating the perception that reconciliation is no longer possible (13). In organizations where people come together for a common purpose, conflicts are natural due to differences in abilities, knowledge, skills, and experience among employees (14). Although conflict is often seen negatively, not all conflicts are destructive. A well-managed conflict can lead to positive outcomes, such as reviewing issues, becoming aware of problems, creating solutions, and improving relationships (15). Unresolved conflict, however, can escalate into mobbing behaviors, and as the intensity of the conflict increases, individuals may begin to experience psychological symptoms. Continuing unresolved issues negatively affect both the physical and psychological well-being of the individual (16).

Conflict styles refer to the behavioral actions that individuals resort to in order to cope with conflict situations (17). In conflict management, the styles developed by Rahim (1985) are defined as "integration," "accommodation," "compromise," "competition," and "avoidance" (14).

This study was conducted to evaluate the extent to which healthcare workers are exposed to mobbing behaviors, to examine whether there is a relationship between the level of mobbing exposure and conflict action styles based on various demographic characteristics of healthcare workers, and to raise awareness about mobbing.

MATERIALS AND METHODS

Type and Purpose of the Study: This study is a descriptive and cross-sectional study conducted to determine the relationship between healthcare workers' exposure to mobbing and their conflict action styles.

Study Location and Period: The research was carried out between May 15, 2023, and August 15, 2023, in healthcare institutions affiliated with the Ministry of Health in Karabük province, including physicians, midwives-nurses, and other healthcare professionals.

Population and Sample of the Study: The study population consisted of physicians, midwives-nurses, and other healthcare personnel (medical secretaries, technicians, laboratory staff, cleaning personnel, physiotherapists) actively working in Karabük and willing to participate in the study. Eleven healthcare personnel who reported experiencing mobbing were excluded from work. Based on a study in the literature examining the relationship between psychological violence and conflict management styles among nurses, the calculated sample size (G-POWER) analysis determined that the minimum sample size required to achieve 80% power and a 95% confidence interval was 246.

Data Collection: The subject, content, and purpose of the study were explained to healthcare workers in written form, and participation was voluntary. The survey was conducted using Google Forms, with data collection tools (scales and survey forms) sent to healthcare workers via a link.

Data Collection Tools: The survey form consisted of three sections. The first section included the Sociodemographic Information Questionnaire, the second section contained the Mobbing Scale, and the third section comprised the Conflict Action Styles Scale.

The Sociodemographic Information Questionnaire included 17 questions about participants' age, gender, education level, marital status, and professional groups.

The Mobbing Scale, in the second section, was developed by Aiello, Deitingner, Nardella, and Bonafede (2008) to measure the exposure of healthcare workers to mobbing. The Turkish version of the Mobbing Scale was validated and tested for reliability by Ayşegül Laleoğlu and Prof. Dr. Emine Özmete in 2013. The Cronbach's alpha internal

consistency coefficient of the survey was 0.948. In this study, the Cronbach's alpha coefficient was found to be $\alpha=0.978$. The statements in the scale were scored on a seven-point Likert scale, ranging from "Strongly Agree - 7" to "Strongly Disagree - 1." A high score on the scale indicated greater exposure to mobbing behaviors, whereas a low score suggested less exposure. Factor analysis results revealed five sub-factors representing mobbing behaviors. The validity and reliability study of the Mobbing Scale by Laleoğlu and Özmete (2013) identified the sub-dimensions as "relationships with colleagues," "threat and harassment," "work and career," "interference with private life," and "commitment to work". The first subdimension consists of 17 items, the second subdimension consists of 7 items, the third subdimension consists of 8 items, the fourth subdimension consists of 4 items, and the fifth subdimension consists of 2 items. The scoring range of the scale is between 38 and 266. The Cronbach's alpha values for these factors were 0.961, 0.904, 0.902, 0.867, and 0.931, respectively.

The Conflict Action Styles Scale, in the third section, was developed by Johnson and Johnson (2008) and was first published in 1981, undergoing multiple revisions over time. The Turkish adaptation of the scale was based on the revision conducted by Prof. Dr. Engin Karadağ and Assoc. Prof. Dr. Ülkü Tosun (2014). The scale consists of 35 attitudinal statements and is divided into five subscales: avoidant, forcing, facilitating, compromising, and oppositional. Each subscale's reliability analysis yielded Cronbach's alpha values of 0.79, 0.76, 0.77, 0.78, and 0.72, respectively. In this study, the Cronbach's alpha values for the sub-dimensions were found to be 0.70, 0.70, 0.75, 0.72, and 0.73. Each subscale was assessed separately, and no total score was calculated. The scores for each subscale ranged from 5 to 35, with higher scores indicating a greater tendency to adopt that particular conflict action style. The scale was designed based on a five-point Likert format, with response options ranging from (1) "I never behave this way" to (5) "I mostly behave this way."

Data Analysis: Statistical analyses were performed using the SPSS 27.0 statistical software. Descriptive statistics such as frequency, percentage, mean, and standard deviation were used. The reliability of the data was assessed using Cronbach's alpha coefficient. The Kolmogorov-Smirnov test was used to determine whether the data followed a normal distribution ($n \geq 30$). Since the Mobbing Scale did not show normal distribution, the Mann-Whitney U test (U-table value) was used to compare the measurement values of two independent groups, and the Kruskal-Wallis test (H-table value) was used for comparisons among three or more groups. Dunnett's test was applied as a post-hoc analysis to determine differences between groups. Since the Conflict Action Styles Scale showed normal distribution, the independent t-test (t-table value) was used to compare two independent groups, and One-Way ANOVA (F-table value) was applied for comparisons among three or more groups. Bonferroni's test was used as a post-hoc analysis to identify group differences. The relationships between the scales were determined using Spearman correlation analysis. All comparisons were evaluated at a 95% confidence interval, with statistical significance set at $p < 0.05$.

Ethical Approval: Research permission was obtained by applying to the Karabuk University Non-

Interventional Clinical Research Ethics Committee, and ethical approval was granted on 08.05.2023 with decision number 2023/1343. Before participating in the study, healthcare workers provided both verbal and written informed consent.

RESULTS

The average age of the 225 healthcare workers who agreed to participate in the study is 32.37 years (30.00 ± 6.839). According to Table 1, 47.5% of the 225 participating healthcare workers are in the 20-29 age group, 65.8% are female, 57.3% are married, and 55.1% have a bachelor's degree. Among the participants, 51.1% are midwives-nurses, while 90.7% work in a training and research hospital. In the study, 69.3% of the participants work in a mixed day-night shift system, 54.2% work 40-47 hours per week, 72.9% are satisfied with their unit, and 59.1% do not regularly smoke (Table 1).

In our study, the total mean score of the healthcare workers on the mobbing scale was determined as 82.15 ± 44.10 (Min: 38; Max: 264). The highest score among the subdimensions of the mobbing scale was found in the "relationships with colleagues" subdimension, with a mean score of 38.99 ± 22.62 (Min: 17; Max: 118).

Table 1: Sociodemographic Characteristics of Healthcare Workers

| | | (n=225) | (%=100) |
|------------------------|-------------------|---------|---------|
| Age | 20-29 | 107 | 47.5 |
| | 30-39 | 72 | 32.0 |
| | 40-49 | 42 | 18.7 |
| | 50 and above | 4 | 1.8 |
| Gender | Female | 148 | 65.8 |
| | Male | 77 | 34.2 |
| Marital Status | Married | 129 | 57.3 |
| | Single | 90 | 40 |
| | Divorced | 6 | 2.7 |
| Education Level | Associate Degree | 40 | 17.8 |
| | Bachelor's Degree | 124 | 55.1 |
| | Postgraduate | 61 | 27.1 |

Table 1: Sociodemographic Characteristics of Healthcare Workers

| | | | |
|----------------------------------|-------------------------------------|-----|------|
| Profession | Midwife-Nurse | 115 | 51.1 |
| | Doctor | 51 | 22.7 |
| | Other Healthcare Worker* | 59 | 26.2 |
| Institution of Employment | Training and Research Hospital | 204 | 90.7 |
| | Family and Community Health Centers | 21 | 9.3 |
| Work Type | Daytime | 69 | 30.7 |
| | Night-Day Mixed | 156 | 69.3 |
| Weekly Working Hours | 40-47 hours | 122 | 54.2 |
| | 47 hours and above | 103 | 45.8 |
| Satisfaction Status | Satisfied | 164 | 72.9 |
| | Not Satisfied | 61 | 27.1 |
| Smoking Status | Smoker | 92 | 40.9 |
| | Non-Smoker | 133 | 59.1 |

*medical secretary, technician, laboratory technician, cleaning staff, physiotherapist

Table 2: Relationship between health workers' perceptions of mobbing and sociodemographic variables

| Variables | Mobbing Scale | Relationships with Colleagues Subdimension | Threat and Harassment Subdimension | Work and Career Obstructions Subdimension | Interference with Private Life Subdimension | Commitment to Work Subdimension |
|--|----------------------|---|---|--|--|--|
| | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ |
| Marital Status | | | | | | |
| Married | 80.000±44.007 | 38.178±22.065 | 11.232±7.899 | 18.697±11.503 | 7.317±5.069 | 4.573±3.166 |
| Single | 87.000±45.002 | 41.155±23.740 | 12.066±7.403 | 20.711±11.855 | 8.522±5.649 | 4.544±2.907 |
| Divorced | 55.666±11.325 | 24.166±7.808 | 7.000±0.000 | 14.000±7.694 | 4.166±0.408 | 3.166±2.206 |
| Test statistics | H=5.171 | H=4.727 | H=9.256 | H=4.001 | H=9.979 | H=0.781 |
| p value | p=0.075 | p=0.094 | p=0.010 | p=0.135 | p=0.007 | p=677 |
| Education Level | | | | | | |
| Associate Degree | 85.352±50.326 | 40.725±25.827 | 12.470±8.882 | 19.352±12.542 | 7.941±6.290 | 4.862±3.492 |
| Bachelor's Degree | 79.373±42.850 | 38.617±22.475 | 10.060±6.980 | 19.373±12.345 | 7.173±5.019 | 4.147±2.896 |
| Postgraduate | 84.796±41.086 | 38.237±20.118 | 13.288±7.251 | 19.406±9.186 | 8.576±4.832 | 5.288±3.023 |
| Test statistics | H=3.132 | H=0.460 | H=24.158 | H=1.716 | H=12.108 | H=10.35 |
| p value | p=0.209 | p=0.795 | p=<0.001 | p=0.424 | p=0.002 | p=0.006 |
| Institution of Employment | | | | | | |
| Training and Research Hospital | 83.725±43.420 | 40.004±22.530 | 11.524±7.414 | 19.740±11.604 | 7.852±5.259 | 4.602±3.080 |
| Family and Community Health Centers | 66.857±48.757 | 29.195±21.611 | 10.761±9.627 | 15.857±11.208 | 6.381±5.607 | 4.666±3.351 |
| Test statistics | U=1220.000 | U=1167.500 | U=1603.000 | U=1567.500 | U=1415.000 | U=2106.500 |
| p value | U=1220.000 | p=<0.001 | p=0.047 | p=0.041 | p=0.008 | p=0.897 |

Table 2: Relationship between health workers' perceptions of mobbing and sociodemographic variables

| Variables | Mobbing Scale | Relationships with Colleagues Subdimension | Threat and Harassment Subdimension | Work and Career Obstructions Subdimension | Interference with Private Life Subdimension | Commitment to Work Subdimension |
|---|-------------------------|--|------------------------------------|---|---|---------------------------------|
| | $\bar{X} \pm \text{SD}$ | $\bar{X} \pm \text{SD}$ | $\bar{X} \pm \text{SD}$ | $\bar{X} \pm \text{SD}$ | $\bar{X} \pm \text{SD}$ | $\bar{X} \pm \text{SD}$ |
| Work Type | | | | | | |
| Daytime | 76.695±46.568 | 34.869±22.173 | 11.173±8.571 | 18.115±11.380 | 7.318±5.561 | 5.217±3.705 |
| Night-Day Mixed | 84.564±42.900 | 40.820±22.645 | 11.576±7.193 | 19.935±11.686 | 7.891±5.185 | 4.339±2.760 |
| Test statistics | U=4353.000 | U=4143.500 | U=4644.500 | U=4730.000 | U=4514.000 | U=4762.500 |
| p value | p=0.022 | p=0.006 | p=0.087 | p=0.144 | p=0.046 | p=0.156 |
| Weekly Working Hours | | | | | | |
| 40-47 hours | 75.303±41.243 | 34.875±19.872 | 10.877±7.631 | 17.352±10.228 | 7.319±5.029 | 4.877±3.402 |
| Over 47 hours | 90.262±46.164 | 43.873±24.715 | 12.135±7.597 | 21.776±12.673 | 8.184±5.586 | 4.291±2.677 |
| Test statistics | U=4840.500 | U=4791.500 | U=5229.500 | U=4946.500 | U=5539.500 | U=5816.500 |
| p value | p=0.003 | p=0.002 | p=0.024 | p=0.006 | p=0.113 | p=0.322 |
| Satisfaction Status | | | | | | |
| Satisfied | 71.250±33.964 | 32.993±16.409 | 10.573±6.241 | 16.286±9.105 | 6.908±4.150 | 4.487±2.985 |
| Not Satisfied | 111.459±54.125 | 55.131±28.558 | 13.819±10.169 | 27.688±13.441 | 9.885±7.169 | 4.934±3.390 |
| Test statistics | U=2348.500 | U=2452.000 | U=3962.000 | U=2161.000 | U=3828.500 | U=4808.500 |
| p value | p=<0.001 | p=<0.001 | p=0.012 | p=<0.001 | p=0.005 | p=0.645 |
| Smoking Status | | | | | | |
| Smoker | 85.272±48.175 | 40.021±23.247 | 12.858±8.887 | 19.510±12.081 | 8.108±5.389 | 4.771±3.298 |
| Non-Smoker | 79.992±41.100 | 38.285±22.236 | 10.481±6.472 | 19.285±11.298 | 7.443±5.235 | 4.496±2.960 |
| Test statistics | U=5489.500 | U=5646.000 | U=4716.000 | U=6057.000 | U=5201.500 | U=5886.000 |
| p value | p=0.190 | p=0.325 | p=0.002 | p=0.898 | p=0.048 | p=0.618 |
| *The group that creates a significant difference between the groups, Kruskal Wallis Test=H, Mann Whitney U Test=U. X: Mean, SD: Standard deviation | | | | | | |

A significant difference was found between the marital status of healthcare workers and the subdimensions of threat and harassment ($p=0.010$) and interference with private life ($p=0.007$). According to the advanced analysis performed to determine the difference, healthcare workers in the divorced group had significantly higher mean scores on the interference with private life subdimension compared to the married or single groups ($p<0.001$). A significant difference was also found between the professional variable and the subdimensions of threat and harassment ($p<0.001$), interference with private life ($p=0.002$), and work commitment ($p=0.006$). Regarding the institution worked at, a significant difference was found between the total score of the mobbing scale ($p=0.001$), the relationships

with coworkers subdimension ($p<0.001$), threat and harassment subdimension ($p=0.047$), job and career-related obstacles subdimension ($p=0.041$), and interference with private life subdimension ($p=0.008$). Additionally, a significant difference was found between satisfaction status and the total score of the mobbing scale ($p<0.001$), relationships with coworkers subdimension ($p<0.001$), threat and harassment subdimension ($p=0.012$), job and career-related obstacles subdimension ($p<0.001$), and interference with private life subdimension ($p=0.005$) (Table 2).

There is no significant difference between healthcare workers' age, gender, and educational status and their perception of mobbing.

Table 3: The Relationship Between Healthcare Workers' Conflict Action Styles and Sociodemographic Variables

| Variables | Avoidant Style Subdimension | Forcing Style Subdimension | Facilitating Style Subdimension | Compromising Style Subdimension | Oppositional Style Subdimension |
|--|-----------------------------|----------------------------|---------------------------------|---------------------------------|---------------------------------|
| | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ | $X \pm SS$ |
| Gender | | | | | |
| Female | 19.121±4.583 | 22.594±4.782 | 24.939±4.776 | 16.358±5.921 | 21.459±3.928 |
| Male* | 21.285±4.895 | 24.220±5.662 | 25.831±5.222 | 17.818±7.677 | 22.675±4.705 |
| Test statistics | t=-3.282 | t=-2.270 | t=-1.287 | t=-2.201 | t=-2.056 |
| p value | p=0.001 | p=0.024 | p=0.200 | p=0.029 | p=0.041 |
| Education Level | | | | | |
| Associate Degree | 21.350±5.051 | 25.075±5.562 | 27.450±4.684 | 17.575±7.605 | 22.825±4.031 |
| Bachelor's Degree | 19.556±4.899 | 22.701±5.002 | 24.766±5.035 | 16.451±6.042 | 21.524±4.381 |
| Postgraduate | 19.508±4.264 | 22.803±4.945 | 24.770±4.576 | 17.213±7.007 | 21.967±4.041 |
| Test statistics | F=2.374 | F=3.483 | F=5.022 | F=0.558 | F=1.449 |
| p value | p=0.095 | p=0.032 | p=0.007 | p=0.573 | p=0.237 |
| Profession | | | | | |
| Doctor | 19.352±4.279 | 21.882±4.563 | 24.549±4.553 | 17.607±7.526 | 21.686±3.916 |
| Midwife-Nurse | 18.791±4.978 | 22.843±5.449 | 24.782±5.418 | 16.173±6.309 | 21.513±4.696 |
| Other Healthcare Workes | 22.389±3.904 | 24.847±4.630 | 26.745±3.941 | 17.542±6.243 | 22.745±3.432 |
| Test statistics | F=12.530 | F=5.144 | F=3.826 | F=4.183 | F=1.725 |
| p value | p=<0.001 | p=0.007 | p=0.023 | p=0.009 | p=0.181 |
| Weekly Working Hours | | | | | |
| 40-47 hours | 20.352±4.695 | 23.795±5.269 | 25.967±4.709 | 16.139±6.409 | 22.327±4.168 |
| Over 47 hours | 19.281±4.865 | 22.388±4.913 | 24.388±5.091 | 17.708±6.739 | 21.339±4.280 |
| Test statistics | F=0.158 | F=0.605 | F=0.326 | F=1.680 | F=0.200 |
| p value | p=0.095 | p=0.041 | p=0.017 | p=0.075 | p=0.082 |
| Satisfaction Status | | | | | |
| Satisfied* | 20.390±4.689 | 23.573±5.163 | 25.939±4.442 | 15.402±5.375 | 22.378±3.885 |
| Not Satisfied | 18.442±4.818 | 22.016±4.964 | 23.377±5.713 | 20.770±7.906 | 20.524±4.853 |
| Test statistics | t=2.749 | t=2.031 | t=3.165 | t=3.776 | t=2.965 |
| p value | p=0.006 | p=0.043 | p=0.002 | p=<0.001 | p=0.003 |
| *The group that creates a significant difference between groups, t = t-test / F = Variance analysis (ANOVA) test | | | | | |

No significant difference was found between the gender variable of healthcare workers and the facilitator style subdimension, while significant differences were found in the other subdimensions. A significant difference was found between the education level of healthcare workers and the facilitator subdimension ($p=0.007$) and the coercive subdimension ($p=0.032$). In advanced analysis, the mean scores of the facilitator subdimension were found to be significantly higher among healthcare

workers with an associate degree compared to those with a bachelor's degree. No significant difference was found in the resistive style subdimension based on the professional variable, but significant differences were found in the other subdimensions. In the advanced analysis, the mean scores of the avoidant subdimension were significantly lower between doctors ($p=0.002$) and midwives-nurses ($p<0.001$) compared to other healthcare workers. According to post-hoc analysis, the mean scores of the coercive

subdimension were significantly lower between other healthcare workers and doctors ($p=0.007$). A significant difference was found between the weekly working hours of healthcare workers and the facilitator subdimension ($p=0.017$) and the coercive subdimension ($p=0.041$). Significant differences were found between all subdimensions and satisfaction status (Table 3).

DISCUSSION

In this study, it was observed that sociodemographic characteristics such as marital status, profession, workplace, work schedule, weekly working hours, job satisfaction, and smoking status influenced perceptions of mobbing behaviors. Participants' marital status affected their perceptions of experiencing mobbing in the "threat and harassment sub-dimension" and the "intervention in private life sub-dimension." Advanced analysis to determine the difference revealed that the mean scores for the intervention in private life sub-dimension were significantly higher in the divorced group compared to the married and single groups. Unlike our study, Akca and colleagues conducted a study on women's perceptions of mobbing in the healthcare sector and found a significant difference in the mobbing experiences of single healthcare workers (18). Similarly, a study conducted between 2016 and 2017 in a state and university hospital in Konya found a significant difference in the levels of mobbing behavior based on marital status, with single healthcare workers experiencing mobbing at a higher rate (19). Differences in findings may be due to factors such as the number of participants, the cultural characteristics of the geographical region, gender distribution, and the type of institution where they work. The higher levels of mobbing behavior experienced by divorced individuals in our study could be related to societal attitudes toward divorce. The perception that divorced healthcare workers may be required to work more shifts due to workload distribution in institutions may have influenced their mobbing experiences.

In this study, a significant difference was found between the professional variable and the threat and harassment sub-dimension, the intervention in private life sub-dimension, and the commitment to work sub-dimension. However, advanced analysis did not reveal any significant differences among specific groups. A study conducted in a state hospital in the Black Sea region among healthcare workers from different departments found that the reputational attack sub-dimension of the mobbing behavior scale varied based on occupational status, with nurses and emergency medical technicians (EMTs) experiencing mobbing more frequently than other employees (20). Another study by Kırılmaz and colleagues in 2015 at Bolu Training and Research Hospital found that risk factors related to mobbing behaviors were higher among physicians (83.3%), nurses (46.9%), and health officers (80%) based on their job titles (21). These differences may be attributed to variations in the number of respondents and sample size. Healthcare workers are considered a high-risk group compared to employees in other service sectors. Issues such as unclear job descriptions, working with insufficient healthcare personnel, heavy workload, stressful environments, and challenging working conditions may contribute to negative behaviors among healthcare professionals.

Our research found a significant difference between the institution where healthcare workers were employed and the mobbing scale, the relationships with colleagues sub-dimension, the threat and harassment sub-dimension, the work and career obstruction sub-dimension, and the intervention in private life sub-dimension, particularly in favor of Training and Research Hospitals. A study conducted by Uysal and colleagues in 2018 at Bolu İzzet Baysal Training and Research Hospital found a significant difference in the level of mobbing experienced by healthcare workers based on their employment affiliation, with those employed under the Ministry of Health experiencing mobbing at a higher rate (22). The similarities between our findings may be due to factors such as the characteristics of inpatient

treatment institutions, bed capacity, insufficient healthcare personnel, high patient volume, and increased workload.

A significant difference was observed between job satisfaction and the total mobbing scale score, the relationships with colleagues sub-dimension, the threat and harassment sub-dimension, the work and career obstruction sub-dimension, and the intervention in private life sub-dimension. In parallel with our study, a study conducted by Yıldırım and Daşbaşı on female social workers employed in the public sector found that participants who were dissatisfied with their institutions had higher mobbing behavior scores (23). Similarities in findings may be attributed to factors such as the work environment, workload, team dynamics, working hours, excessive workload, assignments, and biases.

In this study, it was observed that sociodemographic characteristics such as gender, education level, professional status, weekly working hours, job satisfaction with the unit of employment, and the presence of chronic illness influenced conflict action styles.

It was determined that there was a significant difference in favor of men between the gender variable and the avoiding sub-dimension, the forcing sub-dimension, the compromising sub-dimension, and the resisting sub-dimension. Similar to our study, a study conducted by İkiz and Çatal on teacher candidates found that the forcing sub-dimension of conflict action styles differed significantly in favor of men (24). However, in contrast to our findings, a study conducted by Akpolat and Oğuz in the 2020-2021 academic year in Istanbul with school administrators observed that female administrators used the facilitating, forcing, and avoiding action styles more frequently than male administrators (25). The way men are raised may cause them to exhibit more forceful or resistant behavior when faced with conflicts.

In our study, a significant difference was found between participants' education level and the facilitating and forcing sub-dimensions. In Yılmaz's

study, it was found that among nurses, those in the associate degree and vocational high school groups were more likely to choose the avoidance style in peer conflicts compared to those in the bachelor's and postgraduate groups (26). However, in Özkaya's study on school administrators and teachers in Denizli, no differences were observed in conflict action styles based on education level (27). The differences in research findings may stem from the fact that some studies focus on specific professional groups. Differences in educational level may lead individuals to use various conflict action styles in conflict management.

Our research findings revealed a significant difference in conflict action styles based on professional variables among healthcare workers, specifically in the avoiding, forcing, facilitating, and compromising sub-dimensions. It was found that doctors and midwives/nurses used the avoiding style more frequently than other healthcare workers. Additionally, doctors used the forcing sub-dimension more than other healthcare professionals. A study conducted by Delak and Sirok in 2018 on physicians and nurses in primary healthcare services in Slovenia found that nurses preferred the avoidance style, while physicians preferred the compromising style (28). Similarly, in Akpolat and Oğuz's study on school administrators in Istanbul, assistant principals were found to use the avoiding approach more frequently than school principals (25). These findings are consistent with our results. Professional differences may influence individuals' preferred conflict management styles when dealing with conflicts.

A significant difference was observed in all sub-dimensions of conflict action styles in favor of those who were satisfied with their current situation. Parallel to our findings, a study conducted by Bozkurt and Beydağ in 2023 on nurses working in a private hospital found that 61.2% of participants were satisfied with their workplace, and the resisting and facilitating sub-dimension score averages were high. It is believed that factors such as team motivation, communication among employees,

a high-paced work environment, and personal and cultural differences play a role in the conflict resolution methods used by healthcare workers.

Limitations

The limitations of this study include the fact that it was conducted only with healthcare workers working in the city center of Karabük. Therefore, the findings of this study cannot be generalized to the entire population.

CONCLUSION

In the study, significant statistical results were obtained between some sociodemographic characteristics and various sub-dimensions of both the mobbing perception scale and the conflict action styles scale. It was observed that healthcare workers scored the highest in the “relationships with colleagues” sub-dimension of the mobbing scale, while they scored the highest in the “facilitating approach” sub-dimension of the conflict action styles scale.

It was found that healthcare workers are at significant risk for mobbing, and to prevent this, it is recommended to increase awareness training on mobbing in in-service training programs. Additionally, supportive work environments, management support, fair practices among employees, and the establishment of a mobbing reporting hotline can play an important role in combating mobbing. In general, more comprehensive legal protections should be developed to prevent mobbing behaviors.

REFERENCES

- Güllü C, Çiftçi U. Çalışma yaşamında şiddet ve tacizin önlenmesine ilişkin başvuru rehberi. Ankara: Uluslararası Çalışma Örgütü. 2022.
- Çevirme A. Önlenebilir Bir Halk Sağlığı Sorunu: İşyerlerinde Mobbing ve Sonuçları (The Preventable Public Health Problem: Mobbing and results At Workplaces). *Anadolu Hemşirelik ve Sağlık Bilimleri Dergisi*. 2017;20(3):217-24.
- Cavico FJ, Mujtaba BG, Lawrence E, Muffler SC. Examining the efficacy of the common law tort of intentional infliction of emotional distress and bullying in the context of the employment relationship. *Business Ethics and Leadership*. 2018;2(2):14.
- Yiğit B. Mobbing kavramı: kavramsal bir çerçeve. *Avrasya Sosyal ve Ekonomi Araştırmaları Dergisi*. 2018;5(1):32-42.
- Mujtaba BG, Senathip T. Workplace mobbing and the role of human resources management. 2020.
- Leymann H. The content and development of mobbing at work. *European journal of work and organizational psychology*. 1996;5(2):165-84.
- Tong M, Schwendimann R, Zúñiga F. Mobbing among care workers in nursing homes: A cross-sectional secondary analysis of the Swiss Nursing Homes Human Resources Project. *Int J Nurs Stud*. 2017;66:72-81.
- Şen N. İş hayatında psikolojik şiddet (mobbing): Trabzon örneği. *İmgelem*. 2017;1(1):135-51.
- Acar N, Turunç Ö, Uçku R. Tıp fakültesi uzmanlık öğrencilerinde mobbingin depresyon belirti insidansına etkisi: Bir kohort çalışması. *Turkish Journal of Public Health*. 2020;18(3):195-209.
- Yiğitbaş Ç, Deveci SE. Sağlık çalışanlarına yönelik mobbing. *TTB Mesleki Sağlık ve Güvenlik Dergisi*. 2011;11(42):23-8.
- Sayılan AA, Aydın A. Mobbing among healthcare workers: A review of research findings and methods of struggling. *Journal of Human Sciences*. 2020;17(1):185-93.
- Loretto L, Rosso S, Daga I, Depalmas C, Milia P, Nivoli A. The physician's suicide: study of an Italian sample. *Rivista di psichiatria*. 2020;55(6):9-14.
- Çiğdem A. Okul Yöneticilerinin Çatışma Yönetimi İle İlgili Tezlerin Doküman İncelemesi (2016-2021). *Social Mentality And Researcher Thinkers Journal (Smart Journal)*. 2024;8(64):1824-9.
- Göçer O, Şahin A. Ortaokul müdürlerinin çatışmalara ilişkin görüşleri ve çatışma yönetimi stratejileri. *Kahramanmaraş Sütçü İmam Üniversitesi Sosyal Bilimler Dergisi*. 2022;19(1):44-61.
- Özgüleş B, Çapar H, Boz D, Filiz E, Orhan F, Baş F, Et Al. Sağlık Çalışanlarında Mutsuzluk-Negatif Örgütsel Davranışlar 2023. 216 p.
- Atici Türkeli H. Mobbing ve mobbingin örgütsel bağlılığa etkisi: Sosyal Bilimler Enstitüsü; 2015.
- Karip E. Çatışma Yönetimi. Ankara: Pegem; 2010.
- Akca N, Çakmak A, Şahin H. Cam Tavan Sendromu Ve Mobbing: Kadın Sağlık Çalışanları Üzerine Bir Araştırma. *Akademik Hassasiyetler*. 2022;9(20):89-106.
- Kara S, Kaya ŞD. Sağlık çalışanlarının psikolojik yıldıma ve algılanan sosyal destek düzeylerinin incelenmesi: Konya ilinde bir uygulama. *Selçuk Üniversitesi Sosyal Bilimler Meslek Yüksekokulu Dergisi*. 2021;24(2):381-99.

20. Kılıç T, Çiftçi F, Şener Ş. Sağlık çalışanlarında mobbing ve ilişkili faktörler. 2016.
21. Kırılmaz H, Yorgun S, Atasoy A. Sağlık çalışanlarında psikososyal risk faktörlerini belirlemeye yönelik bir araştırma. Uluslararası Kültürel ve Sosyal Araştırmalar Dergisi. 2016;2(Special Issue 1):66-82.
22. Uysal B, Ekici MA, Önal AC, Kulakoğlu E. Psikolojik Yildırma (Mobing) Ve Çalışan Motivasyonu Arasındaki İlişkinin İncelenmesi. Business & Management Studies: An International Journal. 2019;7(1):280-307.
23. Yıldırım Z, Daşbaşı S. Kamuda Çalışan Kadın Sosyal Çalışmacıların Mobbinge Maruz Kalma Durumlarının İncelenmesi. Toplum ve Sosyal Hizmet. 2022;33(1):179-204.
24. İkiz FE, Çatal U. Öğretmen Adaylarının Çatışma Eylem Stilllerinin Duygusal Öz-Yeterlik Düzeylerine Göre İncelenmesi. Electronic Turkish Studies. 2019;14(4).
25. Akpolat T, Oğuz E. Okul yöneticilerinin analitik ve bütüncül düşünme tarzlarına göre çatışma eylem stillerinin incelenmesi. Yaşadıkça Eğitim. 2022;36(1):35-50.
26. Yılmaz H. Hemşirelerin çatışma yönetim stilleri ve etkileyen faktörlerin incelenmesi: Alanya örneği. 2018.
27. Özkaya MA. Okul yöneticileri ve öğretmenlerin çatışma eylem stillerine ilişkin algılarının belirlenen değişkenlere göre incelenmesi. 2023.
28. Delak B, Şirok K. Physician–nurse conflict resolution styles in primary health care. Nursing open. 2022;9(2):1077-85.